INVASIVE SPECIES FOR HOMEOWNERS

JUMPING WORM

Amynthas agrestis, A. tokioensis, and Metaphire hilgendorfi. Family Megascolecidae

BACKGROUND

Earthworms might be a friendly sight in gardens or your favorite tool for catching fish, but most earthworms in the northeast U.S. are non-native. Jumping worms, a group of species originally from Asia, are invasive species that alter soil qualities and make it inhospitable for some plants and animals. They do this by consuming the

upper organic layer of soil, which leaches nutrients and erodes the ground. This makes it hard for many plants (including garden plants) to grow and threatens even the most well-tended lawns. What's worsehumans spread worms without realizing it, carrying jumping worm egg cases (cocoons) in soil, mulch, potted plants, landscaping equipment, and even the treads of shoes and tires.



▲ Recorded sightings of jumping worm (red) in the Northeast and Midwest U.S. Data from iMapInvasives (2020).

KNOW THE WORM

Life Cycle: Most jumping worms have an annual life cycle. In the spring they hatch from poppy-seed-sized cocoons and after 70-90 days become mature and can produce new cocoons. In the late fall, adults die but the cocoons over-winter to start the next generation.

McCay et al., 2020 luly Most Most juveniles adults Cocoons overwinter in soil until conditions are favorable. Cocoons continue to hatch into the following year. COCOONS

Behavior: These worms thrash wildly and often jump or flip over. Sometimes they cast off the end of their tail.

IN A NUTSHEL

When jumping worms invade, they consume and degrade soil which threatens the future of gardens, forests, and lawns.

Jumping worms are often spread by people through mulch, compost, gardening tools, and treads.

Stop the spread! Use our checklist (pg. 2) to be worm-smart.

Castings: Their castings (feces) appear like coarse coffee grounds that create a loose layer between leaf litter and mineral soil beneath.

UMPING WORM




Look-alikes: Jumping worms might be confused with another invasive worm, the nightcrawler (Lumbricus spp.); both can be large with dark coloring. Jumping worm adults have a smooth milky-white collar (clitellum) close to the head end (14-16 segments away), whereas nightcrawlers have a raised pink-red collar more central along the worm's body.

▲ Jumping worms hold their eggs in cocoons the size of a poppy seed, like this. Can you find the 7 other cocoons?



Jumping Worm Outreach, Research & Management




Note the collar's color and its distance from the head. Credit: UW Madison Arboretum

CHECKLIST

MINIMIZE THE SPREAD

Be a worm-wise buyer









Scout the Soil: Check new mulch, compost, and soil for jumping worms and inquire with providers if measures have been taken to reduce the spread of jumping worms.

Use mulch, compost and soil that are free of jumping worms and cocoons.

If you can't confirm the source is jumping worm-free, only purchase or trade mulch, compost, and soil that has been heated to appropriate temperatures and duration following protocols for reducing pathogens (104 - 130°F for three days is sufficient).







🚳 🔭 Scan the Plants: Check the soil and roots of potted plants and trees for jumping worms or castings before planting them in your yard.


> When the option exists, choose bare-root plants over potted plants, ensuring no soil remains affixed.

> If you find jumping worms in materials you bring in, dispose of all contaminated soil and castings in the trash and kill worms by freezing or leaving in a bag out in the sun, then discard. Alternatively, worms may be killed using vinegar or rubbing alcohol.

KNOW THE SIGNS









present on your property: Focus on preventing spread




present on your property: Focus on monitoring and preventing introductions



before planting to prevent jumping worm introductions

Clean boots, gear, and even roots







When working with logging and landscaping companies, request equipment arrive and leave clear of soil- and encourage your neighbors and local government to do the same.



If jumping worms are present on your property, minimize the sharing and moving of plants where possible. If you do move or share plants, wash roots and share them either bare-root or re-pot in sterile potting soil.



When sharing or moving seedlings and small plants, rinse roots to minimize jumping worm spread when possible: gently massage roots in a basin of water to remove soil clumps, until no soil remains affixed. Afterwards, strain the water and place any solids larger than a poppy seed in a trash bag in the sun before disposing.

TAKE INITIATIVE



Check your property periodically for jumping worms by raking leaf layer, checking underneath.



 Use mustard to count the earthworms on your. property: Mix 1 gallon of water with 1/3 cup of ground yellow mustard seed and pour slowly into the soil. When worms come to the surface, look for jumping worms. Note: 1) this method is safe for most plants 2) mustard is not lethal to earthworms and is not an effective control.



Report jumping worm observations to your state department of natural resources or your local cooperative extension.



Help advance jumping worm research by taking part in citizen science, a bioblitz, or use recording apps like iMapInvasives.

Jumping Worm Outreach, Research, & Management Working Group, 2021

Abby Bezrutczyk, Audrey Bowe, Carrie Brown-Lima, Andrea Dávalos, Annise Dobson, Brad Herrick, Timothy McCay, Kyle Wickings

Questions?

Visit: http://www.nyisri.org/research/jworm-2/

Email: nyisri@cornell.edu

Report your sightings

Go to https://www.imapinvasives.org/ to record any sightings of jumping worms.

We actively use this data for early detection and understanding the invasion.